



Contents lists available at NCBI

## The American Journal of Science and Medical Research

Journal homepage: [www.http://ajsmrjournal.com](http://ajsmrjournal.com)

## Research Article

***In silico* Molecular Docking Studies of compounds from *Rumex vesicarius* against Pancreatic  $\alpha$ -Amylase**Rakesh Davella<sup>1</sup> and Estari Mamidala<sup>2</sup><sup>1,2</sup> Infectious Diseases Research Lab, Department of Zoology, Kakatiya University, Warangal-506009 TS, India.

\*Corresponding author:

E-mail: [drestari@kakatiya.ac.in](mailto:drestari@kakatiya.ac.in)<https://doi.org/10.5281/zenodo.7188942>

Received : 11 October 2019

Accepted; 21 December, 2019

Available online : 28 December, 2019

ISSN: 2377-6196© 2019 The Authors.

<http://ajsmrjournal.com/>

Published by Global Science Publishing Group, USA

**Keywords:** Docking, Pancreatic  $\alpha$ -Amylase, *Rumex vesicarius*, Diabetes mellitus**ABSTRACT**

In spite of the global occurrence of type-2 diabetes mellitus (T2DM) infection and lack of auspicious treatment for Diabetes patients, there are only a few drugs accepted for the managing of infected patients. There is an urgent need to discover newer anti-diabetic drugs with novel mechanism of action and with efforts to reduce attrition rate in early drug discovery stages. The objective of this study is the evaluation of *Rumex vesicarius* compounds for anti-T2DM activity. *In silico* anti-T2DM lead prioritization was performed on a set of known compounds from *R. vesicarius* medicinal plant. The energy minimized structures of these molecules were docked into Pancreatic  $\alpha$ -Amylase. Docking experiments were done using Autodock software for seven compounds docking with  $\alpha$ -Amylase. Rhein was found to be a lead with better docking scores. The results showed that there is scope for the improvement of activity of Isovitexin analogs to discover a potent anti-T2DM compound.

**1. Introduction**

Diabetes mellitus is most common metabolic disease all over the world and number of diabetic patients is still on rise. Diabetes, characterized by hyperglycemia and metabolic disturbance on lipids, carbohydrates, and proteins, affect the life quality of patients by bringing huge pressure to society and public health [1]. Nearly 2.2% of total death in the world is caused by diabetes [2]. Type II diabetes, considered as the common form of diabetes, will affect the health of 8 billion people in the world till 2025 [3]. Persistent hyperglycemia in diabetes mellitus leads to the development of secondary complications including neuropathy, nephropathy, and retinopathy [2]. Type 2 DM (formerly known as non-insulin dependent DM) is the most common form of DM characterized by hyperglycemia, insulin resistance, and relative insulin deficiency [4].

*Rumex vesicarius* Linn. (Polygonaceae) is commonly called as Chukka kura in Telugu, Chukra in Hindi, Bladder Dock in English [5]. *Rumex vesicarius* L. is a wild edible plant used as a sorrel and collected in spring season and eaten fresh or cooked. *Rumex vesicarius* L. has many important medicinal uses such as treatment of hepatic diseases, bad digestion, diuretic, laxative, tonic, analgesic, purgative and antibacterial agents. The plant can be used to reduce biliary disorders and control cholesterol levels [6-7]. Diabetes mellitus is the sixth leading cause of death globally. Many of the drugs have been used in the management of this disease. These drugs have

many side effects and a search for new class of compounds is essential to overcome diabetic problems. Traditionally, a number of plants have been used in various herbal preparations in the management of diabetes and only few of them have been proven scientifically. Anti-bacterial and Antioxidant activities of *Rumex vesicarius* were performed. So far there is no *in silico* anti-diabetic study report so this paper is aimed to report the *in silico* docking of phytochemicals present in this plant against target enzyme,  $\alpha$ -Amylase.

Alpha amylase is an important enzyme which is secreted primarily by the salivary glands and the pancreas contributing its fundamental role in the metabolism of starch and glycogen, which are commonly present in plants, microorganisms and also in higher organisms [8]. Alpha amylase is a target molecule for the treatment of type 2 diabetes mellitus; its inhibitors and its relationship with the disease have been extensively investigated [9-11].

**2. Material and Methods****2.1 Data Set:**

Phytoconstituents from the *R. vesicarius* and standard drug compounds (3D PDB) were downloaded from the IMPPAT database (<https://cb.imsc.res.in/imppat/home>). The molecular properties and structure of the selected compounds (apigenin, Chrysophanol, emodin, Isovitexin, Physcion, thiamine and Vitexin) were shown in table-1.

## 2.2 Bioactive Score Prediction:

Drug score values indicate overall potential of a compound to be a drug candidate. Mol inspiration is a web-based tool used to predict the bioactivity score of the synthesized compounds against regular human receptors such as GPCRs, ion channels, kinases, nuclear receptors, proteases and enzymes

## 2.3 Evaluation of Drug Likelihood:

Lipinski's rule of five is helpful in describing molecular properties of drug compounds required for estimation of important pharmacokinetic parameters such as absorption, distribution, metabolism, excretion and toxicity (ADMET). The rule is helpful in drug design and development. The drug-likeness and molecular property prediction were done by molsoft server (<http://molsoft.com/mprop/>).

## 2.4 Docking Methodology

### 2.4.1 Preparation of Ligand and Protein

The three dimensional structure of the target Pancreatic  $\alpha$ -Amylase (4GQR) was obtained from Protein data bank database (<https://www.rcsb.org/>) and downloaded in PDB format. The ligands and water molecules were removed from the protein and the chemistry of the protein was corrected for the missing hydrogen followed by the energy minimization of the protein. Drug molecule and phytoconstituents optimization, addition of charges and hydrogen bonds was carried out using Autodock tools.

### 2.4.2 Molecular Docking

The docking of 4GQR with selected 40 phytochemical molecules were performed by using Autodock 4 [12]. The docking calculations were verified using docking server. Gasitier partial charges were added to ligand. Nonpolar hydrogen atoms were merged and rotatable hydrogen bonds were defined. Docking calculations were carried out on receptor. Essential hydrogen atoms, kollaman charges and salvation parameters were added affinity (grid) maps 25 Å grid points and 0.500Å were generated using the autogrid program. Autodock parameters set and distance dependent dielectric functions were used in the calculation of the van der Waals and the electrostatic terms, respectively.

Docking simulations were performed using Lamarckian algorithm (LGA) and Solis and Wet local search methods. Initial position torsion and orientation of the drug molecules were set randomly. All rotatable torsions were released during docking. Each docking experiment was derived from 10 different runs that were set to terminate after 250000 energy calculations [13]. The population size was set to 150. During search the translational step 0.2 Å and quaternion and torsion step 5 were applied.

## 3. Results and Discussion

Computational docking is an extremely useful tool to gain an understanding about synthesized compounds and their interactions with biological drug targets, which is very important in drug discovery research. The Molecular Docking software predicted the amino acids in active site region of the studied target proteins.

## 3.1 Evaluation of Drug Likelihood:

Lipinski's rule of five (RO5) is used to evaluate drug likelihood of a chemical compound possessing properties that would make it a likely or potential drug in humans. The oral activity of a drug compound is predicted by calculating certain molecular parameters like log P (partition coefficient), polar surface area, number of hydrogen bond donors, number of hydrogen bond acceptors, and molecular weight. The rule states that most metal complexes with good membrane permeability have log P  $\leq 5$ , number of hydrogen bond acceptors  $\leq 10$ , and number of hydrogen bond donors  $\leq 5$ . In general, an orally active drug has no more than one violation of the given criteria. In the present study, the synthesized ligand and its complexes were found to be in good agreement with the given criteria and can be said to possess good oral bioavailability.

Evaluation of drug likelihood based on Lipinski's rule of five of ligands was showed in Table-1. Based on the drug likelihood evaluation, most of ligands are maintained the Lipinski's rule of five.

## 3.2 Bioactivity Score Prediction:










The pharmacological activity describes the beneficial effects of drugs in living beings. The drug is supposed to bind with a biological target. Biological targets are the most common proteins such as enzymes, ion channels, and receptors. The biological target is also referred to as drug target. The bioactivity scores of the synthesized complexes were calculated for different parameters such as binding to G protein-coupled receptor (GPCR) ligand and nuclear receptor ligand, ion channel modulation, kinase inhibition, amaidotransferase inhibition, and enzyme activity inhibition. All the parameters were calculated with the help of online software Molinspiration ([www.molinspiration.com](http://www.molinspiration.com)), which predicted moderate biological activity for the synthesized complexes. The bioactivity score is given in Table-2. It is known that for metal complexes, if the bioactivity score is more than 0.0, then the complex is active; if it is between -5.0 and 0.0, then the complex is moderately active, and if the bioactivity score is less than -5.0, then it is inactive.












As seen in Table-2, the bioactivity scores of the ligand as well as all the complexes were above -5.0 and 0.0, which clearly indicate that they possess such properties as are enquired for the complexes to act as potential drugs with some modifications in chemical structure.












## 3.3 Docking Interactions:










Physicochemical properties of the 40 compounds of *R.vesicarius* were examined and the results of docking were tabulated. On docking against  $\alpha$ - amylase, the compound Rhein showed greater binding affinity towards enzyme and got a best ligand pose energy of -7.74 and the residues involving in the interaction are: SER:3, PHE:229, PRO:228, ILE:230, ASN:5,250, :GLY:249, LYS:208, 227, TYR:2. In the present study,40 compounds (apigenin, Chrysophanol, emodin, Isovitexin, Physcion and Thiamine, Vitexin etc..) of *Rumex vesicarius* were docked into Pancreatic  $\alpha$ -Amylase enzyme and out of 40, five compounds, Chrysophanol (-7.09 kcal/mol), Isovitexin (-7.31 kcal/mol), Catechin (-7.21 kcal/mol), Rhein (-7.77 kcal/mol), Alloaromadendrene (-7.56 kcal/mol), Methyl dehydroabietate (-7.17 kcal/mol) indicated high binding score (-7.31 kcal/mol).

Table-1. Evaluation of drug likeliness based on Lipinski's rule of five of ligands.

| S.N | Compound Name | Structure   | Molecular formula                                  | Molecular weight | Number of HBA | Number of HBD | Log P |
|-----|---------------|---|--|------------------|---------------|---------------|-------|
| 1   | Apigenin      |    | C <sub>15</sub> H <sub>10</sub> O <sub>5</sub>     | 270.24           | 5             | 3             | 2.46  |
| 2   | Chrysophanol  |    | C <sub>15</sub> H <sub>10</sub> O <sub>4</sub>     | 254.24           | 4             | 2             | 3.54  |
| 3   | emodin        |    | C <sub>15</sub> H <sub>10</sub> O <sub>5</sub>     | 270.24           | 5             | 3             | 3.01  |
| 4   | Isovitexin    |   | C <sub>21</sub> H <sub>20</sub> O <sub>10</sub>    | 432.38           | 10            | 7             | 0.52  |
| 5   | Orientin      |  | C <sub>21</sub> H <sub>20</sub> O <sub>11</sub>    | 448.38           | 11            | 8             | 0.03  |
| 6   | Physcion      |  | C <sub>16</sub> H <sub>12</sub> O <sub>5</sub>     | 284.27           | 5             | 2             | 2.98  |
| 7   | retinol       |  | C <sub>20</sub> H <sub>30</sub> O                  | 286.46           | 1             | 1             | 5.92  |
| 8   | thiamine      |  | C <sub>12</sub> H <sub>17</sub> N <sub>4</sub> O S | 265.36           | 4             | 3             | 0.51  |
| 9   | Vitexin       |  | C <sub>21</sub> H <sub>20</sub> O <sub>10</sub>    | 432.38           | 10            | 7             | 0.52  |

|    |                      |   |  |        |   |   |       |
|----|----------------------|---|--|--------|---|---|-------|
| 10 | Catechin             |    | C <sub>15</sub> H <sub>14</sub> O <sub>6</sub>   | 290.08 | 6 | 5 | 1.88  |
| 11 | Epicatechin          |    | C <sub>15</sub> H <sub>14</sub> O <sub>6</sub>   | 290.08 | 6 | 5 | 1.88  |
| 12 | Iso orientin         |    | C <sub>15</sub> H <sub>15</sub> N O <sub>3</sub> | 257.11 | 3 | 1 | 1.93  |
| 13 | Luteolin             |    | C <sub>15</sub> H <sub>10</sub> O <sub>6</sub>   | 286.05 | 6 | 4 | 2.68  |
| 14 | Quercetin            |    | C <sub>15</sub> H <sub>10</sub> O <sub>7</sub>   | 302.04 | 7 | 5 | 2.11  |
| 15 | Rhein                |  | C <sub>15</sub> H <sub>8</sub> O <sub>6</sub>    | 284.03 | 6 | 3 | 2.19  |
| 16 | 2,6-Dimethylundecane |  | C <sub>13</sub> H <sub>28</sub>                  | 184.22 | 0 | 0 | 6.38  |
| 17 | Docosane             |  | C <sub>22</sub> H <sub>46</sub>                  | 310.36 | 0 | 0 | 10.95 |
| 18 | Dodecane             |  | C <sub>12</sub> H <sub>26</sub>                  | 170.20 | 0 | 0 | 6.13  |
| 19 | Hexacosane           |  | C <sub>26</sub> H <sub>54</sub>                  | 366.42 | 0 | 0 | 12.88 |
| 20 | Eicosane             |  | C <sub>20</sub> H <sub>42</sub>                  | 282.33 | 0 | 0 | 9.98  |

|    |                          |   |            |        |   |   |       |
|----|--------------------------|---|------------|--------|---|---|-------|
| 21 | Tetra methylene_ sulfone |    | C4 H8 O2 S | 120.02 | 2 | 0 | 0.44  |
| 22 | Triacontane              |    | C30 H62    | 422.49 | 0 | 0 | 14.80 |
| 23 | Undecane                 |    | C11 H24    | 156.19 | 0 | 0 | 5.65  |
| 24 | Methyl_ laurate          |    | C13 H26 O2 | 214.19 | 2 | 0 | 5.08  |
| 25 | Methyl_ myristate        |    | C15 H30 O2 | 242.22 | 2 | 0 | 6.04  |
| 26 | methyl_ eicosenate       |   | C21 H40 O2 | 324.30 | 2 | 0 | 9.08  |
| 27 | methyl_ pentadecanoate   |  | C16 H32 O2 | 256.24 | 2 | 0 | 6.52  |
| 28 | Methyl_ palmitate        |  | C17 H34 O2 | 270.26 | 2 | 0 | 7.00  |
| 29 | methyl_ margarate        |  | C18 H36 O2 | 284.27 | 2 | 0 | 7.49  |
| 30 | methyl_ linoleate        |  | C19 H34 O2 | 294.26 | 2 | 0 | 7.08  |
| 31 | Methyl_ stearate         |  | C19 H38 O2 | 298.29 | 2 | 0 | 7.97  |

|    |                        |   |            |        |   |   |       |
|----|------------------------|---|------------|--------|---|---|-------|
| 32 | Methyl_eicosanoate     |    | C22 H44 O2 | 340.33 | 2 | 0 | 9.25  |
| 33 | Methyl_dehydroabietate |    | C21 H30 O2 | 314.22 | 2 | 0 | 5.66  |
| 34 | Methyl_docosanoate     |    | C23 H46 O2 | 354.35 | 2 | 0 | 9.90  |
| 35 | Methyl_lignocerate     |    | C25 H50 O2 | 382.38 | 2 | 0 | 10.86 |
| 36 | Alloaromadendrene      |    | C15 H24    | 204.19 | 0 | 0 | 4.85  |
| 37 | Beta-Elemene           |   | C15 H24    | 204.19 | 0 | 0 | 5.39  |
| 38 | Cadina-1_4-diene       |  | C15 H24    | 204.19 | 0 | 0 | 6.02  |
| 39 | Cis-Limonene           |  | C10 H16 O  | 152.12 | 1 | 0 | 2.95  |
| 40 | Humulene               |  | C15 H24    | 204.19 | 0 | 0 | 5.47  |

The former compound is thus an effective inhibitor among the other compounds that can stop the function of  $\alpha$ -amylase. However, further *in vitro* and *in vivo* studies of individual phytoconstituents is needed to validate their biological potential. Ball and socket model of respective drug molecule and phytoconstituents interacting with active site are shown in [Table-3](#).

#### 4. Conclusion

Understanding the interactions between proteins and ligands is crucial for the pharmaceutical and functional food industries. The emergence of bioinformatics has offered a platform to explore diseases at molecular level using computational tools. The Protein-Ligand interaction plays a significant role in structure based drug designing. Finally from this analysis it was found that, among the all compounds six compounds are effectively inhibit T2DM-  $\alpha$ -amylase and the phytochemicals of this plant can act as T2DM-  $\alpha$ -amylase inhibitors.




**Table-2. Bioactivity score of the ligands**

| S. No. | Compound               | Parameters of Bioactivity score |                       |                  |                         |                    |                  |
|--------|------------------------|---------------------------------|-----------------------|------------------|-------------------------|--------------------|------------------|
|        |                        | GPCR ligand                     | Ion channel modulator | Kinase inhibitor | Nuclear receptor ligand | Protease inhibitor | Enzyme inhibitor |
| 1      | Apigenin               | -0.07                           | -0.09                 | 0.18             | 0.34                    | -0.25              | 0.26             |
| 2      | Chrysophanol           | -0.23                           | -0.17                 | 0.06             | 0.02                    | -0.26              | 0.16             |
| 3      | Emodin                 | -0.14                           | -0.14                 | 0.07             | 0.17                    | -0.21              | 0.21             |
| 4      | Isovitexin             | 0.12                            | 0.02                  | 0.15             | 0.23                    | 0.04               | 0.47             |
| 5      | Orientin               | 0.12                            | -0.14                 | 0.20             | 0.20                    | 0.01               | 0.45             |
| 6      | Physcion               | -0.17                           | -0.23                 | 0.04             | 0.11                    | -0.23              | 0.14             |
| 7      | Retinol                | -0.01                           | 0.32                  | -0.25            | 1.02                    | -0.16              | 0.66             |
| 8      | Thiamine               | 0.26                            | 0.01                  | -0.37            | -1.72                   | -0.64              | 1.12             |
| 9      | Vitexin                | 0.13                            | -0.14                 | 0.19             | 0.23                    | 0.03               | 0.46             |
| 10     | catechin               | 0.41                            | 0.14                  | 0.09             | 0.60                    | 0.26               | 0.47             |
| 11     | Epicatechin            | 0.41                            | 0.14                  | 0.09             | 0.60                    | 0.26               | 0.47             |
| 12     | Isoorientin            | 0.11                            | 0.01                  | 0.16             | 0.20                    | 0.01               | 0.46             |
| 13     | Luteolin               | -0.02                           | -0.07                 | 0.26             | 0.39                    | -0.22              | 0.28             |
| 14     | Quercetin              | -0.06                           | -0.19                 | 0.28             | 0.36                    | -0.25              | 0.28             |
| 15     | Rhein                  | -0.08                           | -0.10                 | 0.01             | 0.29                    | -0.06              | 0.28             |
| 16     | 2_6-Dimethylundecane   | -0.57                           | -0.18                 | -0.93            | -0.67                   | -0.56              | -0.27            |
| 17     | Docosane               | 0.02                            | 0.00                  | -0.07            | 0.02                    | -0.02              | 0.03             |
| 18     | Dodecane               | -0.71                           | -0.23                 | -0.93            | -0.83                   | -0.85              | -0.35            |
| 19     | Hexacosane             | 0.04                            | 0.00                  | -0.04            | 0.04                    | 0.04               | 0.03             |
| 20     | Icosane                | -0.04                           | 0.00                  | -0.14            | -0.05                   | -0.11              | 0.03             |
| 21     | tetramethylene_sulfone | -3.73                           | -3.88                 | -3.91            | -3.92                   | -3.56              | -3.67            |
| 22     | Triacontane            | 0.04                            | 0.00                  | -0.04            | 0.04                    | 0.03               | 0.02             |
| 23     | Undecane               | -0.84                           | -0.31                 | -1.08            | -0.98                   | -0.98              | -0.44            |
| 24     | alloaromadendrene      | -0.67                           | -0.47                 | -0.98            | -0.21                   | -0.67              | -0.30            |
| 25     | beta-elemene           | -0.36                           | 0.18                  | -1.02            | 0.43                    | -0.38              | 0.30             |
| 26     | Cadina-1_4-diene       | -0.33                           | -0.15                 | -0.85            | -0.09                   | -0.77              | 0.19             |
| 27     | cis- -Limonene_oxide   | -0.65                           | -0.40                 | -2.04            | 0.15                    | -0.63              | 0.37             |
| 28     | Humulene               | -0.14                           | 0.02                  | -0.93            | 0.34                    | -0.67              | 0.31             |
| 29     | Methyl_laurate         | -0.41                           | -0.13                 | -0.73            | -0.43                   | -0.46              | -0.11            |
| 30     | Methyl_myristate       | -0.24                           | -0.07                 | -0.51            | -0.24                   | -0.28              | -0.02            |
| 31     | methyl_eicosenate      | 0.02                            | -0.08                 | -0.23            | 0.12                    | 0.07               | 0.07             |
| 32     | methyl_pentadecanoate  | -0.17                           | -0.06                 | -0.42            | -0.16                   | -0.20              | 0.01             |
| 33     | Methyl_palmitate       | -0.11                           | -0.05                 | -0.34            | -0.09                   | -0.13              | 0.04             |
| 34     | methyl_margarate       | -0.07                           | -0.05                 | -0.28            | -0.04                   | -0.08              | 0.05             |
| 35     | methyl_linoleate       | 0.15                            | 0.07                  | -0.20            | 0.14                    | 0.03               | 0.23             |
| 36     | Methyl_stearate        | -0.03                           | -0.04                 | -0.23            | 0.00                    | -0.03              | 0.05             |
| 37     | methyl_eicosanoate     | 0.03                            | -0.02                 | -0.20            | 0.07                    | 0.14               | 0.04             |
| 38     | methyl_dehydroabietate | 0.40                            | 0.20                  | -0.26            | 0.85                    | 0.02               | 0.36             |
| 39     | methyl_docosanoate     | 0.02                            | -0.04                 | -0.16            | 0.05                    | 0.06               | 0.04             |
| 40     | methyl_lignocerate     | 0.02                            | -0.04                 | -0.15            | 0.05                    | 0.06               | 0.04             |


Table-3. Phytochemical compounds docked against T2DM- Pancreatic  $\alpha$ - amylase

| Sl.No | Plant Name              | Compound Name          | Binding energy | Inhibition constant | Intermolecular energy |
|-------|-------------------------|------------------------|----------------|---------------------|-----------------------|
| 1     | <i>Rumex vesicarius</i> | Apigenin               | -6.68 kcal/mol | 12.60 uM            | -7.88 kcal/mol        |
| 2     |                         | Chrysophanol           | -7.09 kcal/mol | 6.34 uM             | -7.69 kcal/mol        |
| 3     |                         | Emodin                 | -6.92 kcal/mol | 8.52 uM             | -7.81 kcal/mol        |
| 4     |                         | Isovitexin             | -7.31 kcal/mol | 4.35 uM             | -10.30 kcal/mol       |
| 5     |                         | Physcion               | -6.93 kcal/mol | 8.38 uM             | -7.82 kcal/mol        |
| 6     |                         | Retinol                | -7.54 kcal/mol | 2.96 uM             | -9.33 kcal/mol        |
| 7     |                         | Thiamine               | -6.09 kcal/mol | 34.23 uM            | -7.88 kcal/mol        |
| 8     |                         | Vitexin                | -6.48 kcal/mol | 17.68 uM            | -9.47 kcal/mol        |
| 9     |                         | Catechin               | -7.21 kcal/mol | 5.14 uM             | -9.00 kcal/mol        |
| 10    |                         | Epicatechin            | -6.58 kcal/mol | 15.11 uM            | -8.37 kcal/mol        |
| 11    |                         | Isoorientin            | -6.24 kcal/mol | 26.67 uM            | -9.52 kcal/mol        |
| 12    |                         | Luteolin               | -6.80 kcal/mol | 10.42 uM            | -8.29 kcal/mol        |
| 13    |                         | Quercetin              | -6.59 kcal/mol | 14.87 uM            | -8.38 kcal/mol        |
| 14    |                         | Rhein                  | -7.74 kcal/mol | 2.14 uM             | -8.93 kcal/mol        |
| 15    |                         | 2,6-Dimethylundecane   | -5.13 kcal/mol | 173.59 uM           | -7.52 kcal/mol        |
| 16    |                         | Docosane               | -5.61 kcal/mol | 76.90 uM            | -11.28 kcal/mol       |
| 17    |                         | Dodecane               | -4.50 kcal/mol | 506.68 uM           | -7.18 kcal/mol        |
| 18    |                         | Hexacosane             | -4.85 kcal/mol | 280.01 uM           | -11.71 kcal/mol       |
| 19    |                         | Icosane                | -4.76 kcal/mol | 322.90 uM           | -9.83 kcal/mol        |
| 20    |                         | Tetramethylene_Sulfone | -4.28 kcal/mol | 728.99 uM           | -4.28 kcal/mol        |
| 21    |                         | Triacontane            | -4.87 kcal/mol | 267.49 uM           | -12.93 kcal/mol       |
| 22    |                         | Undecane               | -4.24 kcal/mol | 777.24 uM           | -6.63 kcal/mol        |
| 23    |                         | Alloaromadendrene      | -7.56 kcal/mol | 2.87 uM             | -7.56 kcal/mol        |
| 24    |                         | Beta-Elemene           | -6.97 kcal/mol | 7.77 uM             | -7.87 kcal/mol        |
| 25    |                         | Cadina-1_4-Diene       | -7.17 kcal/mol | 5.53 uM             | -7.47 kcal/mol        |
| 26    |                         | Cis-_-Limonene_Oxide   | -5.63 kcal/mol | 74.83 uM            | -5.93 kcal/mol        |
| 27    |                         | Humulene               | -6.96 kcal/mol | 7.89 uM             | -6.96 kcal/mol        |
| 28    |                         | Methyl_Laurate         | -4.68 kcal/mol | 369.45 uM           | -7.96 kcal/mol        |
| 29    |                         | Methyl_Myristate       | -4.30 kcal/mol | 700.39 uM           | -8.18 kcal/mol        |
| 30    |                         | Methyl_Eicosenate      | -4.62 kcal/mol | 409.12 uM           | -10.29 kcal/mol       |
| 31    |                         | Methyl_Pentadecanoate  | -4.74 kcal/mol | 333.16 uM           | -8.92 kcal/mol        |
| 32    |                         | Methyl_Palmitate       | -4.59 kcal/mol | 432.60 uM           | -9.06 kcal/mol        |
| 33    |                         | Methyl_Margarate       | -4.65 kcal/mol | 393.04 uM           | -9.42 kcal/mol        |
| 34    |                         | Methyl_Linoleate       | -5.36 kcal/mol | 118.44 uM           | -9.83 kcal/mol        |
| 35    |                         | Methyl_Stearate        | -4.36 kcal/mol | 633.76 uM           | -9.43 kcal/mol        |
| 36    |                         | Methyl_Eicosanoate     | -4.26 kcal/mol | 755.02 uM           | -9.93 kcal/mol        |
| 37    |                         | Methyl_Dehydroabietate | -7.17 kcal/mol | 1.69 uM             | -8.77 kcal/mol        |
| 38    |                         | Methyl_Docosanoate     | -4.58 kcal/mol | 436.35 uM           | -10.85 kcal/mol       |
| 39    |                         | Methyl_Lignocerate     | -4.96 kcal/mol | 230.67 uM           | -11.82 kcal/mol       |
| 40    |                         | Orientin               | -5.31 kcal/mol | 4.35 uM             | -10.30 kcal/mol       |
| 41    | REFERENCE               | Acarbose               | -2.24 kcal/mol | 22.98 mM            | -8.80 kcal/mol        |


Figure-1. *In silico* Molecular Docking Studies of compounds (a- Chrysophanol; b- Isovitexin; c- Catechin; d- Rhein; e- Alloaromadendrene; f- methyl\_dehydroabietate) from *Rumex vesicarius* against Pancreatic  $\alpha$ -Amylase









(b)




(c)



(d)



(e)



(f)

Therefore, this study emphasizes the importance of small molecules from various plant sources and their use to enhance protein-ligand interaction studies in silico. Further investigations can be done on our in silico approach to produce more effective and potential T2DM-  $\alpha$ -amylase inhibitors through ligand based drug designing approaches..

#### Acknowledgement:

To authors thank to the Department of Zoology, Kakatiya University, Warangal, TS, India for giving permission to carry out this work.

#### Conflicting Interests

The authors have declared that no conflicting interests exist.

#### References

- [1]. S. Ramachandran, A. Rajasekaran, and N. Adhirajan, "In vivo and in vitro antidiabetic activity of Terminalia paniculata bark: an evaluation of possible phytoconstituents and mechanisms for blood glucose control in diabetes," *ISRN Pharmacology*, vol. 2013, Article ID 484675, 10 pages, 2013.
- [2]. B. A. Sheikh, L. Pari, A. Rathinam, and R. Chandramohan, "Trans-anethole, a terpenoid ameliorates hyperglycemia by regulating key enzymes of carbohydrate metabolism in streptozotocin induced diabetic rats," *Biochimie*, vol. 112, pp. 57–65, 2015.
- [3]. G. Winkler, T. Hidvegi, G. Vandrofi, S. Balogh, and G. Jermendy, "Risk-stratified screening for type 2 diabetes in adult subjects: results from Hungary," *Diabetologia*, vol. 54, p. S119, 2011.
- [4]. Maitra A, Abbas AK. Endocrine system. In: Kumar V, Fausto N, Abbas AK (eds). *Robbins and Cotran Pathologic basis of disease* (7th ed) 2005. Philadelphia, Saunders; 1156-1226.

- [5]. Madhavashetty K Dr, Shivaji K, Tulasirao K, Flowering plants of Chittoor district, Andhra Pradesh, India; 298.
- [6]. Elegami, A.A.; Almagboul, A.Z.; El faith, M.A.O. and El Tohami, M.S.: Sudanese plants used in folkloric medicine: Screening for antibacterial activity. Part X. *Fitoterapia* 2001; 72: 810-817.
- [7]. Al-Quran, S.: Ethnopharmacological survey of wild medicinal plants in Showbak, Jordan. *Journal of Ethnopharmacology* 2009; 123 : 45-50.
- [8]. Kandra, L.  $\alpha$ -Amylases of medical and industrial importance. *J. Mol. Struct. THEOCHEM*, 2003, 666, 487-498.
- [9]. Nickavar, B., and Abolhasani, L. (2013) Bioactivity-guided separation of an  $\alpha$ -amylase inhibitor flavonoid from *Salvia virgata*. *Iran. J. Pharm. Res.* 12, 57-61
- [10]. Yadav, R., Bhartiya, J. P., Verma, S. K., and Nandkeoliar, M. K. (2013) The evaluation of serum amylase in the patients of type 2 diabetes mellitus, with a possible correlation with the pancreatic functions. *J. Clin. Diagn. Res.* 7, 1291-1294
- [11]. Ozdemir S, et al. Isolation of a Novel Thermophilic *Anoxybacillus flavithermus* SO-13, Production, Characterization and Industrial Applications of its Thermostable Alpha-Amylase. *J Bioprocess Biotech.* 2015; 5: 237.
- [12]. Bikadi Z and Hazai E: Application of the PM6 semi-empirical method to modeling proteins enhances docking accuracy of AutoDock. *Journal of Cheminformatics* 2009; 1: 15.
- [13]. Halgren TA: Merck molecular force field. I. Basis, form, scope, parametrization, and performance of MMFF94. *Journal of Computational Chemistry* 1998.